

Purified anti-eIF4E

Catalog # / Size: 606902 / 200 µl (20 Western blots)

Clone: Poly6069

Isotype: Rabbit IgG

Immunogen: Recombinant (partial), N-terminal

Reactivity: Human

Preparation: The antibody was purified by antigen-affinity chromatography.


Formulation: This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 50% glycerol.

Storage: Upon receipt, store frozen at -20°C.

Applications:

Applications: WB - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by Western blotting. Western blotting, suggested working dilution(s): Use 10 µl per 5 ml antibody dilution buffer for each mini-gel. It is recommended that the reagent be titrated for optimal performance for each application.



MCF-7 cell extract was resolved by electrophoresis, transferred to nitrocellulose, and probed with rabbit anti-eIF4E antibody. Proteins were visualized using a donkey anti-rabbit secondary conjugated to HRP and a chemiluminescence detection system.

Description: eIF4E (also known as eukaryotic translation initiation factor 4E) is a 25 kD member of the eukaryotic initiation factor 4E family. This ubiquitously expressed cytoplasmic protein recognizes and binds 7-methylguanosine containing mRNA "CAP" during initiation of protein synthesis. eIF4E facilitates ribosome binding by inducing unwinding of mRNA secondary structures. Activity of this protein is inhibited by binding of 4E-BP and PRH; expression is increased by H₂O₂ treatment. Activity can also be regulated by integrin α₆β₄ ligation. Phosphorylation increases binding to mRNA CAPs and formation of eIF4F complex. eIF4E forms a trimeric complex with eIF4G, eIF4A (called eIF4F). Also associates with 4E-BP and PRH. The Poly6069 antibody has recognized human eIF4E and has been shown to be useful for Western blotting.

Other Names: Eukaryotic translation initiation factor 4E

Antigen References:

1. Rychlik W, *et al.* 1987. *P. Natl. Acad. Sci. USA* 84:945.
2. Topisirovic I, *et al.* 2003. *EMBO J.* 22:689.
3. Shenberger J, *et al.* 2002. *Am. J. Respir. Cell Mol. Biol.* 27:250.

Related Products:

Product
Purified anti-eIF2α

Purified anti-eIF2γ

HRP Donkey anti-rabbit IgG (minimal x-reactivity)

Clone

Poly6067

Poly6068

Poly4064

Application

WB

WB

ELISA, IHC, WB



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.