

Recombinant Human GM-CSF (ELISA Std.)

Catalog # / Size: 572409 / 4 pack

Source: Human GM-CSF, amino acids Ala18- Glu144 (Accession# AAA52578) was expressed in *E. coli*, approximately 14.6 kD.

Preparation: ELISA Standard 4-Pack contains four vials of recombinant human GM-CSF at >5 ng per vial. Recommended for ELISA application only. Standards are calibrated against a BioLegend Master Calibrator.

Formulation: Lyophilized in sterile-filtered PBS, pH 7.2, containing 1% BSA.

Storage: Upon receipt, store unopened vials between 2°C and 8°C immediately and use within 12 months from date of receipt. Prior to use, reconstitute the lyophilized powder with 0.2 ml of PBS containing a carrier protein (e.g., 1% BSA, protease free), pH7.4. Re-cap vial, vortex. Allow the reconstituted standard to sit at room temperature for 15 minutes, vortex again to mix completely. The reconstituted standard stock solution can be aliquoted into polypropylene vials and stored at -70°C for up to one month. Do not re-use diluted standards. Use a manual defrost freezer and avoid repeated freeze thaw cycles.

Applications:

Applications: ELISA

Recommended Usage: Each lot of this protein is quality control tested by ELISA assay. For use as an ELISA standard, a standard curve comprised of doubling dilutions from 500 pg/ml to 4 pg/ml is suggested. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: This GM-CSF protein is useful as a standard for a human GM-CSF sandwich ELISA, using unlabelled anti-GM-CSF antibody, clone BVD2-23B6 (catalog #502202) for capture and biotinylated BVD2-21C11 (catalog #502304) antibody for detection.

Description: GM-CSF plays a key role in signaling emergency hemopoiesis (predominantly myelopoiesis) in response to infection, including the production of granulocytes and macrophages in the bone marrow and their maintenance, survival, and functional activation at sites of injury or insult. The receptor for GM-CSF is a heterodimer that comprises a major binding subunit (GMR α) and a major signaling subunit (β c). The receptor subunits are always coexpressed on the surface of leukocytes, with β c being expressed at lower levels than GMR α . Certain nonhemopoietic cell types have also been reported to express the GM-CSF receptor and to respond to GM-CSF stimulation *in vitro*, although the *in vivo* significance of these observations remains uncertain.

Antigen References:

1. Hercus TR, *et al.* 2009 *Blood* 114:1289-1298.
2. Hayashida K, *et al.* 1990 *Proc Natl Acad Sci USA*. 87:9655-9659.
3. Walker F, *et al.* 1985 *EMBO J* 4:933-939.
4. Kitamura T, *et al.* 1989 *J. Cell Physiol.* 140:323.

Related Products:

Product
Purified anti-human GM-CSF
Biotin anti-human GM-CSF

Clone
BVD2-23B6
BVD2-21C11

Application
ELISA Capture, IP, WB
ELISA Detection, ELISPOT
Detection, ICFC
ELISA
ELISA
ELISA

Human GM-CSF ELISA MAX™ Standard
Human GM-CSF ELISA MAX™ Deluxe
LEGEND MAX™ Human GM-CSF ELISA Kit with
Pre-coated Plates



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.