

# Product Data Sheet

## Human TruStain FcX™ (Fc Receptor Blocking Solution)

**Catalog # / Size:** 422301 / 50 tests  
422302 / 200 tests

**Storage:** Store between 2°C and 8°C.

### Applications:


**Applications:** FC - *Quality tested*  
ICFC, ICC - *Reported in the literature*

**Recommended Usage:** It is recommended to add 5 µl of Human TruStain FcX™ per 100 µl of cell suspension, mix and incubate at room temperature for 5-10 minutes prior to staining with antibody of interest. It is not necessary to wash cells between these blocking and immunostaining steps.


**Application Notes:** This buffer contains specialized human IgG. It is not recommended to be used for staining human IgG. Handle as biohazard agent.

**Application References:**

1. de la Cruz J, *et al.* 2011. *J. Immunol.* 186:2282. PubMed
2. Luo L, *et al.* 2011. *Exp Mol Pathol.* 91:596. PubMed
3. Szerlip NJ, *et al.* 2012. *PNAS.* 109:3041. PubMed
4. Bromley SK, *et al.* 2013. *J. Immunol.* 190:970. PubMed
5. Trujillo G, *et al.* 2013. *J. Immunol.* 191:848. PubMed
6. Edin S, *et al.* 2013. *PLoS One.* 18:74982. PubMed
7. Euzebio Alves VT, *et al.* 2013. *Infect Immun.* 81:4399. PubMed
8. Borg ZD, *et al.* 2014. *J. Immunol.* 192:189. PubMed
9. Klingenberg R, *et al.* 2014. *Eur Heart J.* PubMed
10. Lines JL, *et al.* 2014. *Cancer Res.* 74:1924. PubMed
11. Decker D, *et al.* 2014. *Lupus.* 23:802. PubMed
12. Jitschin R, *et al.* 2014. *Blood.* 123:2663. PubMed
13. Anderson ME Jr, *et al.* 2014. *Otolaryngol Head Neck Surg.* 150:880. PubMed
14. Giurisato E, *et al.* 2014. *Mol Biol Cell.* 25:1769. PubMed
15. Shi J, *et al.* 2014. *Blood.* 123:4015. PubMed
16. Kusner LL, *et al.* 2014. *PLoS One.* 9:102231. PubMed
17. Fujisawa D, *et al.* 2014. *J Allergy Clin Immunol.* 134:622. PubMed
18. Bhardwaj N, *et al.* 2014. *J Virol.* 88:11108. PubMed
19. Lieber S, *et al.* 2014. *Mol Pharmacol.* 87:162. PubMed
20. Olganier D, *et al.* 2014. *PLoS Pathog.* 10:1004566. PubMed



Human TruStain FcX™ treated (blue) or non-treated (red) THP-1 cells stained with an irrelevant mouse IgG1 PE mAb (above). Dotted histogram represents unstained cells.



Human TruStain FcX™ treated (filled histograms) or non-treated (open histograms) THP-1 cells stained with anti-human HLA-DR PE (red) or an isotype control (IgG2a PE, blue) mAbs. Note: non-treated cells show false-negative HLA-DR staining due to high binding of mouse IgG2a isotype.

**Description:** Human Fc receptors (FcRs) are expressed on a variety of cells, such as monocytes, granulocytes, B cells and dendritic cells. The cells with FcR expression sometimes give false positive or false negative results of immunofluorescent staining due to the FcRs-mediated Ig Fc binding. Human TruStain FcX™ is specially formulated for blocking the FcR-involved unwanted staining without interfering with antibody-mediated specific staining of human cells. Human TruStain FcX™ is compatible with flow cytometric staining with anti-human CD16 (clone 3G8), CD32 (clone FUN-2), and CD64 (clone 10.1) antibodies.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.