

Cell-Vive™ GMP Ultra-LEAF™ Purified anti-human CD56 (NCAM) SF Antibody

Catalog# / Size	362571 / 100 µg 362572 / 1 mg
Clone	5.1H11
Other Names	Leu-19, NKH1
Isotype	Mouse IgG1, κ
Description	<p>CD56 is a single transmembrane glycoprotein also known as NCAM (neural cell adhesion molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is expressed on NK and NKT cells. CD56 is also expressed in the brain (cerebellum and cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparan sulfate.</p> <p>Cell-Vive™ GMP Ultra-LEAF™ Purified anti-human CD56 (NCAM) SF Antibody was GMP manufactured under serum-free conditions including serum-free hybridoma cell culture, without additional animal or human derived materials or preservatives. This antibody contains ultra-low levels of endotoxin (<0.01EU/µg of protein), is filtered through a 0.1 µm membrane, and is tested negative for mycoplasma and microbial growths.</p> <p>Cell-Vive™ GMP Ultra-LEAF™ Purified anti-human CD56 (NCAM) SF Antibody is for research and further ex vivo bioprocessing use only.</p>

Product Details

Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	0.1 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is < 0.01 EU/µg of the protein (< 0.001 ng/µg of the protein) as determined by the LAL test.
Preparation	The Ultra-LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.
Concentration	1.0 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. This Ultra-LEAF™ solution contains no preservative; handle under aseptic conditions.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	Cell-Vive™ GMP Ultra-LEAF™ Purified anti-human CD56 (NCAM) SF Antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.1 µm filtered) is recommended for highly sensitive assays. The antibody was GMP manufactured under serum-free conditions including serum-free hybridoma cell culture, without additional animal or human-derived materials or preservatives. This antibody is intended for flow cytometry and for research or further ex vivo bioprocessing use only.
Application References	<ol style="list-style-type: none"> Walsh FS, <i>et al.</i> 1981. <i>Nature</i>. 289:60. (FC) Pavlath GK, <i>et al.</i> 1986. <i>J. Cell Biol.</i> 102:124. (FC) Pavlath GK, <i>et al.</i> 1989. <i>Nature</i>. 337:570. (FC) Pulido R, <i>et al.</i> 1988. <i>J. Immunol.</i> 140:3851. (FC)
(PubMed link indicates BioLegend citation)	

Antigen Details

Structure	Ig superfamily, single transmembrane or GPI-anchored glycoprotein
Distribution	NK cells, T subset, neural tissue, some LGL and myeloid leukemias
Function	Adhesion
Ligand/Receptor	Heparan sulfate
Antigen References	<ol style="list-style-type: none"> 1. Lanier L, <i>et al.</i> 1991. <i>J. Immunol.</i> 146:4421-6. 2. Hemperly J, <i>et al.</i> 1990. <i>J. Mol. Neurosci.</i> 2:71-8. 3. Cremer H, <i>et al.</i> 1994. <i>Nature.</i> 367:455-9.
Gene ID	4684


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD56 (NCAM), APC anti-human CD56 (NCAM), PerCP/Cyanine5.5 anti-human CD56 (NCAM), PE anti-human CD56 (NCAM), PE/Cyanine7 anti-human CD56 (NCAM), APC/Cyanine7 anti-human CD56 (NCAM), PE/Cyanine5 anti-human CD56 (NCAM), Alexa Fluor® 647 anti-human CD56 (NCAM), Alexa Fluor® 488 anti-human CD56 (NCAM), Pacific Blue™ anti-human CD56 (NCAM), Alexa Fluor® 700 anti-human CD56 (NCAM), PerCP anti-human CD56 (NCAM), Brilliant Violet 650™ anti-human CD56 (NCAM), Brilliant Violet 510™ anti-human CD56 (NCAM), Brilliant Violet 421™ anti-human CD56 (NCAM), Biotin anti-human CD56 (NCAM), Brilliant Violet 605™ anti-human CD56 (NCAM), Brilliant Violet 570™ anti-human CD56 (NCAM), Brilliant Violet 711™ anti-human CD56 (NCAM), PE/Dazzle™ 594 anti-human CD56 (NCAM), FITC anti-human CD56 (NCAM), Brilliant Violet 785™ anti-human CD56 (NCAM), Ultra-LEAF™ Purified anti-human CD56 (NCAM), APC/Fire™ 750 anti-human CD56 (NCAM), Brilliant Violet 750™ anti-human CD56 (NCAM), TotalSeq™-A0047 anti-human CD56 (NCAM), TotalSeq™-C0047 anti-human CD56 (NCAM), TotalSeq™-B0047 anti-human CD56 (NCAM), Spark NIR™ 685 anti-human CD56 (NCAM), KIRAVIA Blue 520™ anti-human CD56 (NCAM), GMP PE anti-human CD56 (NCAM), TotalSeq™-D0047 anti-human CD56 (NCAM), GMP APC anti-human CD56 (NCAM)

Product Data



Human peripheral blood lymphocytes were stained with Cell-Vive™ GMP Ultra-LEAF™ anti-human CD56 (NCAM) SF antibody (Clone 5.1H11, filled histogram) or isotype control (Clone MOPC-21, open histogram) followed by FITC anti-mouse IgG stain.

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587