

## Biotin anti-human $\beta$ 2-microglobulin Antibody

<b>Catalog# / Size</b>	316308 / 100 $\mu$ g
<b>Clone</b>	2M2
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	$\beta$ 2M, $\beta$ 2-M, beta2-microglobulin b2-M, b2M
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	$\beta$ 2-microglobulin ( $\beta$ 2M) is a 12 kD nonpolymorphic Ig like protein. It is a non-membrane-anchored glycoprotein and is noncovalently associated with 39-44 kD polymorphic heavy chains of MHC class I molecules to form HLA class I antigen complex. In association with HLA class I, $\beta$ 2M is expressed on all leukocytes, platelets, endothelial cells, and epithelial cells. $\beta$ 2M plays an essential role both in governing MHC class I molecules stability and in promoting antigen binding and presenting the antigen to CD3/TCR complex of CD8 <sup>+</sup> T cells.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	African Green, Baboon, Cynomolgus, Rhesus, Pig
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Purified human $\beta$ 2-microglobulin
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is $\leq 0.5$ $\mu$ g per million cells in 100 $\mu$ l volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: Western blotting, and ELISA.
<b>Application References</b>	1. Meissner TB, <i>et al.</i> 2010. <i>Proc Natl Acad Sci USA</i> . <a href="#">PubMed</a> 2. Rizvi SM, <i>et al.</i> 2011. <i>J. Immunol.</i> 186:2309. <a href="#">PubMed</a> 3. Meissner TB, <i>et al.</i> 2012. <i>J Immunol.</i> 188:4951. <a href="#">PubMed</a> .
<b>(PubMed link indicates BioLegend citation)</b>	
<b>Product Citations</b>	1. Tischer DK <i>et al.</i> 2019. eLife. 8 pii: e42498. <a href="#">PubMed</a> 2. Mendoza JL, <i>et al.</i> 2020. eLife. 9:00. <a href="#">PubMed</a>
<b>RRID</b>	AB_493689 (BioLegend Cat. No. 316308)

### Antigen Details

---

<b>Structure</b>	Nonpolymorphic Ig like structure, noncovalently associated with heavy chain of MHC class I molecules, 12 kD
------------------	---

<b>Distribution</b>	Leukocytes, platelets, endothelial cells, epithelial cells
<b>Function</b>	Govern MHC class I molecule stability, promote MHC class I molecules binding and presenting peptide antigens to CD8 <sup>+</sup> T cells
<b>Ligand/Receptor</b>	CD3/TCR complex, CD8
<b>Cell Type</b>	Endothelial cells, Epithelial cells, Leukocytes, Platelets
<b>Biology Area</b>	Immunology
<b>Molecular Family</b>	MHC Antigens
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Engelhard VH. 1994. <i>Curr. Opin. Immunol.</i> 6:13.</li> <li>2. Williams DB, et al. 1989. <i>J. Immunol.</i> 142:2796.</li> <li>3. Danliczyk UG and TL. Delovitch. 1994. <i>J. Immunol.</i> 153:3533.</li> <li>4. Williams A, et al. 2002. <i>Tissue Antigens</i> 59:3.</li> </ol>
<b>Gene ID</b>	<a href="#">567</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-human  $\beta$ 2-microglobulin, FITC anti-human  $\beta$ 2-microglobulin, PE anti-human  $\beta$ 2-microglobulin, Biotin anti-human  $\beta$ 2-microglobulin, APC anti-human  $\beta$ 2-microglobulin, HRP anti-human  $\beta$ 2-microglobulin, PE/Cyanine7 anti-human  $\beta$ 2-microglobulin, APC/Fire™ 750 anti-human  $\beta$ 2-microglobulin, PerCP/Cyanine5.5 anti-human  $\beta$ 2-microglobulin, PE/Dazzle™ 594 anti-human  $\beta$ 2-microglobulin, TotalSeq™-A0057 anti-human  $\beta$ 2-microglobulin, TotalSeq™-C0057 anti-human  $\beta$ 2-microglobulin, TotalSeq™-B0057 anti-human  $\beta$ 2-microglobulin

## Product Data



Human peripheral blood lymphocytes stained with biotinylated 2M2, followed by Sav-PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587