

Alexa Fluor[®] 700 anti-mouse CD8a Antibody

Catalog# / Size	100729 / 25 µg 100730 / 100 µg
Clone	53-6.7
Regulatory Status	RUO
Other Names	T8, Lyt2, Ly-2
Isotype	Rat IgG2a, κ
Description	CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form the α(CD8a)/β(CD8b) heterodimer and α/α homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8 α/β heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR α/β T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8 α/α homodimer is expressed on a subset of γ/δ TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase lck.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Mouse thymus or spleen
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor [®] 700 under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The suggested use of this reagent is ≤ 0.25 µg per 10⁶ cells in 100 µl volume. It is highly recommended that the reagent be titrated for optimal performance for each application.</p> <p>* Alexa Fluor[®] 700 has a maximum emission of 719 nm when it is excited at 633nm / 635nm. Prior to using Alexa Fluor[®] 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.</p> <p>Alexa Fluor[®] and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p>View full statement regarding label licenses</p>
Excitation Laser	Red Laser (633 nm)
Application Notes	Clone 53-6.7 antibody competes with clone 5H10-1 antibody for binding to thymocytes ³ . The 53-6.7 antibody has been reported to block antigen presentation via MHC class I and inhibit T cell responses to IL-2. This antibody has also been used for depletion of CD8a ⁺ cells. Additional reported applications (for the relevant formats) include: immunoprecipitation ^{1,3} , <i>in vivo</i> and <i>in vitro</i> cell depletion ^{2,10,15} , inhibition of CD8 T cell proliferation ³ , blocking of cytotoxicity ^{3,4} , immunohistochemical staining ^{5,6} of acetone-fixed frozen sections and zinc-fixed paraffin-embedded sections, and spatial biology (IBEX) ^{29,30} . Clone 53-6.7 is not recommended for immunohistochemistry of formalin-fixed paraffin sections. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays or in

vivo studies (Cat No. 100746).

Application References

(PubMed link indicates
BioLegend citation)

1. Ledbetter JA, et al. 1979. *Immunol. Rev.* 47:63. (IHC, IP)
2. Hathcock KS. 1991. *Current Protocols in Immunology.* 3.4.1. (Deplete)
3. Takahashi K, et al. 1992. *P. Natl. Acad. Sci. USA* 89:5557. (Block, IP)
4. Ledbetter JA, et al. 1981. *J. Exp. Med.* 153:1503. (Block)
5. Hata H, et al. 2004. *J. Clin. Invest.* 114:582. (IHC)
6. Fan WY, et al. 2001. *Exp. Biol. Med.* 226:1045. (IHC)
7. Shih FF, et al. 2006. *J. Immunol.* 176:3438. (FC)
8. Kamimura D, et al. 2006. *J. Immunol.* 177:306.
9. Bouwer HGA, et al. 2006. *P. Natl. Acad. Sci. USA* 103:5102. (FC, Deplete)
10. Kao C, et al. 2005. *Int. Immunol.* 17:1607. [PubMed](#)
11. Ko SY, et al. 2005. *J. Immunol.* 175:3309. (FC) [PubMed](#)
12. Rasmussen JW, et al. 2006. *Infect. Immun.* 74:6590. [PubMed](#)
13. Lee CH, et al. 2009. *Clin. Cancer Res.* [PubMed](#)
14. Geiben-Lynn R, et al. 2008. *Blood* 112:4585. (Deplete) [PubMed](#)
15. Kingeter LM, et al. 2008. *J. Immunol.* 181:6244. [PubMed](#)
16. Guo Y, et al. 2008. *Blood* 112:480. [PubMed](#)
17. Andrews DM, et al. 2008. *J. Virol.* 82:4931. [PubMed](#)
18. Britschqui MR, et al. 2008. *J. Immunol.* 181:7681. [PubMed](#)
19. Kenna TJ, et al. 2008. *Blood* 111:2091. [PubMed](#)
20. Jordan JM, et al. 2008. *Infect. Immun.* 76:3717. [PubMed](#)
21. Todd DJ, et al. 2009. *J. Exp. Med.* 206:2151. [PubMed](#)
22. Bankoti J, et al. 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
23. Medyouf H, et al. 2010. *Blood* 115:1175. [PubMed](#)
24. Riedl P, et al. 2009. *J. Immunol.* 183:370. [PubMed](#)
25. Apte SH, et al. 2010. *J. Immunol.* 185:998. [PubMed](#)
26. Bankoti J, et al. 2010. *Toxicol. Sci.* 115:422. (FC) [PubMed](#)
27. del Rio ML, et al. 2011. *Transpl. Int.* 24:501. (FC) [PubMed](#)
28. Cui L, et al. 2015. *J Control Release.* 206:220. [PubMed](#)
29. Radtke AJ, et al. 2020. *Proc Natl Acad Sci U S A.* 117:33455-65. (SB) [PubMed](#)
30. Radtke AJ, et al. 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

Product Citations

1. Adams NM et al. 2018. *Immunity.* 48(6):1172-1182 . [PubMed](#)
2. Ding L et al. 2018. *Cell reports.* 25(11):2972-2980 . [PubMed](#)
3. Sadtler K, et al. 2017. *Tissue Eng Part A.* 23:1044. [PubMed](#)
4. Zhang YS, et al. 2018. *Cancer Biol Ther.* 19:735. [PubMed](#)
5. Zhu X et al. 2019. *Cell reports.* 27(4):1205-1220 . [PubMed](#)
6. Yue X, et al. 2019. *Nat Commun.* 10:2011. [PubMed](#)
7. Qi X, et al. 2019. *Nat Commun.* 10:2141. [PubMed](#)
8. Li C, et al. 2018. *Cell.* 174:285. [PubMed](#)
9. Wang F, et al. 2018. *Carcinogenesis.* 39:889. [PubMed](#)
10. Celis-Gutierrez J et al. 2019. *Cell Rep.* 27(11):3315-3330 . [PubMed](#)
11. He W et al. 2018. *Immunity.* 49(6):1175-1190 . [PubMed](#)
12. Jackstadt R, et al. 2019. *Cancer Cell.* 36:319. [PubMed](#)
13. Ji L et al. 2019. *Immunity.* 51(2):272-284 . [PubMed](#)
14. Christopher S Garris et al. 2018. *Immunity.* 49(6):1148-1161 . [PubMed](#)
15. Ying Zhang et al. 2017. *Cancer cell.* 32(3):377-391 . [PubMed](#)
16. Kim MY, et al. 2017. *Plant Biotechnol J.* 1.729166667. [PubMed](#)
17. Zhang H, et al. 2019. *Mol Cell.* 76:110. [PubMed](#)
18. Kim C, et al. 2019. *Cell Rep.* 29:2202. [PubMed](#)
19. Yoshida H, et al. 2019. *Cell.* 176:897. [PubMed](#)
20. Zhang D, et al. 2020. *Signal Transduct Target Ther.* 5:24. [PubMed](#)
21. Annu K, et al. 2020. *Sci Rep.* 10:2359. [PubMed](#)
22. Qiao J, et al. 2019. *Cancer Cell.* 35:901. [PubMed](#)
23. Bronisz-Budzynska I, et al. 2019. *Skelet Muscle.* 9:22. [PubMed](#)
24. Alexander RK, et al. 2020. *eLife.* 9:e54090.. [PubMed](#)
25. Shimizu K, et al. 2020. *Molecular Cell.* 77(5):937-950.e6.. [PubMed](#)
26. Resch TK, et al. 2018. *Sci Rep.* 0.722916667. [PubMed](#)
27. Diaz-Salazar C, et al. 2020. *Cell Rep.* 32:108186. [PubMed](#)
28. Engelbertsen D, et al. 2013. *Arterioscler Thromb Vasc Biol.* 33:637. [PubMed](#)
29. McPhee C, et al. 2013. *J Immunol.* 191:4581. [PubMed](#)
30. Bunn P, et al. 2014. *J Immunol.* 192:3709. [PubMed](#)
31. Carty S, et al. 2014. *PLoS One.* 9:106659. [PubMed](#)
32. Herz J, et al. 2015. *J Exp Med.* 212: 1153 - 1169. [PubMed](#)
33. Montes de Oca M, et al. 2016. *PLoS Pathog.* 12: 1005398. [PubMed](#)
34. Solaymani-Mohammadi S, et al. 2016. *J Leukoc Biol.* 99: 475 - 482. [PubMed](#)
35. Ma C, et al. 2016. *Nature.* 531:253-257. [PubMed](#)
36. Engel I, et al. 2016. *Nat Immunol.* 10.1038/ni.3437. [PubMed](#)
37. Moodley D, et al. 2016. *Proc Natl Acad Sci U S A.* 113: 9852 - 9857. [PubMed](#)
38. Rosnagl S, et al. 2016. *PLoS Biol.* 14: 1002562. [PubMed](#)
39. Pérol L, et al. 2016. *Nat Commun.* 7:13027. [PubMed](#)
40. Panciera T, et al. 2016. *Cell Stem Cell.* 19:725-737. [PubMed](#)
41. Cerina M, et al. 2017. *Brain Behav Immun.* 59:103-117. [PubMed](#)
42. Nakagawa P, et al. 2017. *Physiol Rep.* 10.14814/phy2.13084. [PubMed](#)
43. Wang X, et al. 2020. *Nat Commun.* 4.505555556. [PubMed](#)
44. Fang H, et al. 2020. *Nat Commun.* 4.661805556. [PubMed](#)
45. Lewandowski JP, et al. 2020. *Genome Biol.* 1.039583333. [PubMed](#)

46. Gruber T, *et al.* 2020. JCI Insight. 5:00. [PubMed](#)
47. Zhang D, *et al.* 2020. Signal Transduct Target Ther. 5:24. [PubMed](#)
48. Mender I, *et al.* 2020. Cancer Cell. 38(3):400-411.e6. [PubMed](#)
49. Hamaidi I, *et al.* 2020. Cell Metabolism. 32(3):420-436.e12. [PubMed](#)
50. Hall JA, *et al.* 2020. Cell Metabolism. 32(4):665-675.e6. [PubMed](#)
51. Starkl P, *et al.* 2020. Immunity. 53(4):793-804.e9. [PubMed](#)
52. Biancur DE, *et al.* 2020. Cell Metabolism. 33(1):199-210.e8. [PubMed](#)
53. He Y, *et al.* 2021. Cell Metabolism. 33(5):988-1000.e7. [PubMed](#)
54. Dallari S, *et al.* 2021. Cell Host Microbe. 29(6):1014-1029.e8. [PubMed](#)

RRID AB_493702 (BioLegend Cat. No. 100729)
 AB_493703 (BioLegend Cat. No. 100730)

Antigen Details

Structure	Ig superfamily, CD8 α chain, 34 kD
Distribution	Most thymocytes, T cell subset, some NK cells, lymphoid dendritic cells
Function	Co-receptor for TCR
Ligand/Receptor	MHC class I molecule
Antigen References	<ol style="list-style-type: none"> 1. Barclay A, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press. 2. Zamoyska R. 1994. <i>Immunity</i> 1:243. 3. Ellmeier W, <i>et al.</i> 1999. <i>Annu. Rev. Immunol.</i> 17:523.
Gene ID	12525


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse CD8a, Biotin anti-mouse CD8a, FITC anti-mouse CD8a, PE anti-mouse CD8a, PE/Cyanine5 anti-mouse CD8a, Purified anti-mouse CD8a, PE/Cyanine7 anti-mouse CD8a, APC/Cyanine7 anti-mouse CD8a, Alexa Fluor® 488 anti-mouse CD8a, Alexa Fluor® 647 anti-mouse CD8a, Pacific Blue™ anti-mouse CD8a, Alexa Fluor® 700 anti-mouse CD8a, PerCP/Cyanine5.5 anti-mouse CD8a, PerCP anti-mouse CD8a, Brilliant Violet 421™ anti-mouse CD8a, Brilliant Violet 570™ anti-mouse CD8a, Brilliant Violet 650™ anti-mouse CD8a, Brilliant Violet 605™ anti-mouse CD8a, Ultra-LEAF™ Purified anti-mouse CD8a, Brilliant Violet 711™ anti-mouse CD8a, Brilliant Violet 785™ anti-mouse CD8a, Brilliant Violet 510™ anti-mouse CD8a, Purified anti-mouse CD8a (Maxpar® Ready), Alexa Fluor® 594 anti-mouse CD8a, PE/Dazzle™ 594 anti-mouse CD8a, APC/Fire™ 750 anti-mouse CD8a, GoInVivo™ Purified anti-mouse CD8a, TotalSeq™-A0002 anti-mouse CD8a, Spark Blue™ 550 anti-mouse CD8a, Spark NIR™ 685 anti-mouse CD8a, TotalSeq™-C0002 anti-mouse CD8a, TotalSeq™-B0002 anti-mouse CD8a, Spark YG™ 570 anti-mouse CD8a, PE/Fire™ 640 anti-mouse CD8a, PE/Fire™ 700 anti-mouse CD8a, Spark Blue™ 574 anti-mouse CD8a Antibody, Spark Violet™ 423 anti-mouse CD8a Antibody, Spark UV™ 387 anti-mouse CD8a

Product Data



C57BL/6 mouse splenocytes stained with CD8 (clone 53-6.7) Alexa Fluor® 700 (filled histogram) or rat IgG2a, κ Alexa Fluor® 700 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587