

## Purified anti-mouse CD45.2 (Maxpar<sup>®</sup> Ready) Antibody

<b>Catalog# / Size</b>	109843 / 100 µg
<b>Clone</b>	104
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Ly-5.2, LCA
<b>Isotype</b>	Mouse (SJL) IgG2a, κ
<b>Description</b>	CD45.2 is an alloantigen of CD45, expressed by Ly5.2 bearing mouse strains (e.g., A, AKR, BALB/c, CBA/Ca, CBA/J, C3H/He, C57BL, C57BR, C57L, C58, DBA/1, DBA/2, NZB, SWR, 129). CD45, a member of the protein tyrosine phosphatase (PTP) family, is a 180-240 kD glycoprotein expressed on all hematopoietic cells except mature erythrocytes and platelets. There are multiple isoforms in the mouse that play key roles in TCR and BCR signal transduction. These isoforms are very specific to the activation and maturation states of the cell as well as specific cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.

### Product Details

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	B10.S mouse thymocytes and splenocytes
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and EDTA.
<b>Preparation</b>	The antibody was purified by affinity chromatography.
<b>Concentration</b>	1.0 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C.
<b>Application</b>	<a href="#">FC - Quality tested</a> <a href="#">CyTOF<sup>®</sup> - Verified</a>
<b>Recommended Usage</b>	This product is suitable for use with the <a href="#">Maxpar<sup>®</sup> Metal Labeling Kits</a> . For metal labeling using Maxpar <sup>®</sup> Ready antibodies, proceed directly to the step to Partially Reduce the Antibody by adding 100 µl of Maxpar <sup>®</sup> Ready antibody to 100 µl of 4 mM TCEP-R in a 50 kDa filter and continue with the protocol. Always refer to the latest version of Maxpar <sup>®</sup> User Guide when conjugating Maxpar <sup>®</sup> Ready antibodies.
<b>Application Notes</b>	The 104 antibody does not react with mouse cells expressing the CD45.1 alloantigen. Additional reported applications (for the relevant formats) include: immunoprecipitation <sup>4</sup> , <i>in vivo</i> and <i>in vitro</i> blocking of B cell responses <sup>1,2</sup> , and immunohistochemical staining of acetone-fixed frozen sections <sup>3</sup> .
<b>Additional Product Notes</b>	Maxpar <sup>®</sup> is a registered trademark of Standard BioTools Inc.
<b>Application References</b>	<ol style="list-style-type: none"> <li>1. Yakura H, <i>et al.</i> 1983. <i>J. Exp. Med.</i> 157:1077. (Block)</li> <li>2. Yakura H, <i>et al.</i> 1986. <i>J. Immunol.</i> 136:2729. (Block)</li> <li>3. Suzuki K, <i>et al.</i> 2000. <i>Immunity</i> 13:691. (IHC)</li> <li>4. Shen FW, <i>et al.</i> 1986. <i>Immunogenetics</i> 24:146. (IP)</li> <li>5. Baldwin TA and Hogquist KA. 2007. <i>J. Immunol.</i> 179:837.</li> <li>6. Pascal V, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:1751.</li> <li>7. Burman AC, <i>et al.</i> 2007. <i>Blood</i> 110:1064.</li> <li>8. Kincaid EZ, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:3187.</li> <li>9. Phan TG, <i>et al.</i> 2007. <i>Nature Immunol.</i> 8:992.</li> <li>10. Nakano-Yokomizo T, <i>et al.</i> 2011. <i>J. Exp. Med.</i> 208:1661. <a href="#">PubMed</a></li> <li>11. Wen T, <i>et al.</i> 2013. <i>PNAS.</i> 110:6067. <a href="#">PubMed</a></li> <li>12. Kohlmeier JE, <i>et al.</i> 2008. <i>Immunity.</i> 29:101. (FC) <a href="#">PubMed</a></li> </ol>

RRID

AB\_2563751 (BioLegend Cat. No. 109843)

## Antigen Details

<b>Structure</b>	Protein tyrosine phosphatase (PTP) family, 180-240 kD
<b>Distribution</b>	All hematopoietic cells except mature erythrocytes and platelets of the CD45.2 strain of mice
<b>Function</b>	Phosphatase, T and B cell activation
<b>Ligand/Receptor</b>	Galectin-1, CD2, CD3, CD4
<b>Biology Area</b>	Cell Biology, Immunology, Inhibitory Molecules, Innate Immunity, Neuroscience, Neuroscience Cell Markers
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	1. Suzuki K, <i>et al.</i> 2000. <i>Immunity</i> 13:691.
<b>Gene ID</b>	<a href="#">19264</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Biotin anti-mouse CD45.2, FITC anti-mouse CD45.2, PE anti-mouse CD45.2, Purified anti-mouse CD45.2, APC anti-mouse CD45.2, Alexa Fluor® 488 anti-mouse CD45.2, Alexa Fluor® 647 anti-mouse CD45.2, Pacific Blue™ anti-mouse CD45.2, Alexa Fluor® 700 anti-mouse CD45.2, APC/Cyanine7 anti-mouse CD45.2, PerCP anti-mouse CD45.2, PerCP/Cyanine5.5 anti-mouse CD45.2, PE/Cyanine7 anti-mouse CD45.2, Brilliant Violet 421™ anti-mouse CD45.2, Brilliant Violet 570™ anti-mouse CD45.2, Brilliant Violet 650™ anti-mouse CD45.2, Brilliant Violet 510™ anti-mouse CD45.2, Brilliant Violet 785™ anti-mouse CD45.2, Brilliant Violet 605™ anti-mouse CD45.2, Purified anti-mouse CD45.2 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse CD45.2, Brilliant Violet 711™ anti-mouse CD45.2, Alexa Fluor® 594 anti-mouse CD45.2, APC/Fire™ 750 anti-mouse CD45.2, TotalSeq™-A0157 anti-mouse CD45.2, TotalSeq™-B0157 anti-mouse CD45.2, TotalSeq™-C0157 anti-mouse CD45.2, Brilliant Violet 750™ anti-mouse CD45.2, Spark Blue™ 550 anti-mouse CD45.2, Spark NIR™ 685 anti-mouse CD45.2

## Product Data



C57BL/6 mouse bone marrow cells stained with 147Sm-anti-CD45.2 (104). Data provided by DVS Sciences.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

