

Alexa Fluor® 647 anti-H2A.X Antibody

Catalog# / Size	600205 / 25 µg 600206 / 100 µg
Clone	W16171A
Regulatory Status	RUO
Other Names	H2A.X Variant Histone, H2A Histone Family Member X, Histone H2A.X, Histone H2AX, H2AFX
Isotype	Rat IgG2a, κ
Description	Histone subunit H2A, along with subunits 2B, 3, and 4, make up the core subunits of the nucleosome octamer. An octamer contains two protomers of each subunit tightly wrapped around a ~147 bp segment of DNA. Histones have integral roles in chromatin integrity, genomic stability, and gene regulation. Post-translational modification of histones in response to certain stimuli results in alterations of nucleosomal positioning relative to DNA. Histone H2A.X is a non-allelic variant of Histone 2A that harbors a C-terminal extension and is essential for checkpoint mediated cell cycle arrest and DNA double-stranded break (DSB) repair in response to both endogenous and exogenous agents, as well as meiotic recombination events and immunoglobulin class switching in lymphocytes. Phosphorylation of C-terminal residue serine 139 by ATM (γ-H2A.X) results in the recruitment of DSB-repair machinery. Phosphorylation of H2A.X is also critical for chromatin fragmentation during apoptosis.

Product Details

Verified Reactivity	Human, Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Synthetic human histone H2A.X peptide (127-143) conjugated to KLH.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunocytochemistry. For immunocytochemistry, a concentration range of 1.0 - 5.0 µg/mL is recommended. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.</p> <p>Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p>View full statement regarding label licenses</p>
Excitation Laser	Red Laser (633 nm)
Application Notes	<p>This product is a monoclonal antibody raised against the C-terminus of H2A.X (residues 127-143); BioLegend's existing antibody against H2A.X (Poly6133, cat# 613302) is a polyclonal antibody which was generated against (partial), N-terminal H2A.X.</p> <p>This clone is not recommended for ChIP (Chromatin Immunoprecipitation) assays (as determined by in-house testing).</p>
RRID	<p>AB_2820097 (BioLegend Cat. No. 600205)</p> <p>AB_2832825 (BioLegend Cat. No. 600206)</p>

Antigen Details

Structure	Histone H2A.X is a 143 amino acid protein with a predicted molecular weight of 15.1 kD.
Distribution	Ubiquitous tissue expression; nuclear localization
Function	H2A.X, upon phosphorylation, promotes DNA repair and maintains genomic stability. Important for recombination between immunoglobulin switch regions.
Interaction	ATM, MDC1, TP53BP1, BRCA1, MRE11, RAD50, NBN
Biology Area	Apoptosis/Tumor Suppressors/Cell Death, Cell Biology, Cell Cycle/DNA Replication, Chromatin Remodeling/Epigenetics, DNA Repair/Replication

Antigen References

1. Chen CC, *et al.* 2017. *Proc. Natl. Acad. Sci.* 114: 7665.
2. Natale F, *et al.* 2017. *Nat. Commun.* 8: 15760.
3. Bhargava R, *et al.* 2017. *Proc. Natl. Acad. Sci.* 114: 728.
4. Weyemi U, *et al.* 2016. *Nat. Commun.* 7: 10711.
5. Rezaeian AH, *et al.* 2017. *Nat. Cell. Biol.* 19: 38.
6. Horn S, *et al.* 2015. *Biochim. Biophys. Acta.* 1853: 2199.
7. Reina-San-Martin B, *et al.* 2003. *J. Exp. Med.* 197:1767
8. Celeste A, *et al.* 2002. *Science* 296:922.
9. Mannironi C, *et al.* 1989. *Nucleic Acids Res.* 17:9113.

Gene ID [3014](#)


Related Protocols

[Immunocytochemistry Staining Protocol](#)

Other Formats

Purified anti-H2A.X, Alexa Fluor® 594 anti-H2A.X, Alexa Fluor® 647 anti-H2A.X, Alexa Fluor® 488 anti-H2A.X

Product Data



HeLa cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with Triton X-100 for 10 minutes, and blocked with 5% FBS for 60 minutes. Cells were then intracellularly stained with 1 µg/mL (1:500 dilution) of either Alexa Fluor® 647 Rat IgG2a, κ Isotype Ctrl Antibody (Cat. No. 400526, panel A) or Alexa Fluor® 647 anti-H2A.X antibody (panel B) overnight at 4°C. Nuclei were counterstained with DAPI and the image was captured with a 60X objective.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

