

APC/Fire™ 750 anti-human CD66b Recombinant Antibody

Catalog# / Size	396907 / 25 tests 396908 / 100 tests
Clone	QA17A51
Regulatory Status	RUO
Other Names	CD67, CGM6, NCA-95, CEACAM8
Isotype	Mouse IgG1, κ
Description	CD66b is a 95-100 kD glycosylphosphatidylinositol (GPI)-linked protein also known as CD67, CGM6, and NCA-95. CD66b is a member of the immunoglobulin superfamily, carcinoembryonic antigen (CEA)-like subfamily. CD66b, expressed on granulocytes, has been reported to induce activation in neutrophils and to be involved in heterophilic adhesion with CD66c.

Product Details

Verified Reactivity	Human
Antibody Type	Recombinant
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.
Excitation Laser	Red Laser (633 nm)
RRID	AB_2820054 (BioLegend Cat. No. 396907) AB_2820055 (BioLegend Cat. No. 396908)

Antigen Details

Distribution	Granulocytes/neutrophils
Function	Cell adhesion, neutrophil activation
Ligand/Receptor	CD66b
Cell Type	Granulocytes
Biology Area	Immunology
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	

1. Kuijpers T, *et al.* 1993. *J. Immunol.* 151:4934
2. Kuroki M, *et al.* 1992. *J. Leuk. Biol.* 52:551

Gene ID [1088](#)


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-human CD66b Recombinant Antibody, Purified anti-human CD66b Recombinant Antibody, APC/Fire™ 750 anti-human CD66b Recombinant Antibody, PE/Dazzle™ 594 anti-human CD66b Recombinant Antibody, PerCP/Cyanine5.5 anti-human CD66b Recombinant Antibody, PE/Cyanine7 anti-human CD66b Recombinant Antibody, APC anti-human CD66b Recombinant Antibody, FITC anti-human CD66b Recombinant Antibody

Product Data



Human peripheral blood was stained with True-Stain Monocyte Blocker™ (Cat. No. 426102), CD15 FITC and CD66b (clone QA17A51) APC/Fire™ 750 (left) or, Mouse IgG1, κ APC/Fire™ 750 isotype control (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587