

Purified anti-PSD95 Antibody (Previously Covance catalog# MMS-5183)

Catalog# / Size	810302 / 25 µL 810301 / 100 µL
Clone	K28/74
Regulatory Status	RUO
Other Names	Disks large homolog 4, discs large homolog 4, Tax interaction protein 15, synapse-associated protein 90, postsynaptic density protein 95, post-synaptic density protein 95
Previously	Covance Catalog# MMS-5183
Isotype	Mouse IgG1, κ
Description	PSD95, also known as SAP-90, is a member of the MAGUK family, encoded by the the DLG4 gene. At postsynaptic sites, it mediates the scaffolding of NMDA receptors, AMPA receptors, potassium channels, and other associated signaling protein clusters. It plays an important role in synaptic plasticity and the stabilization of synaptic changes during long-term potentiation.

Product Details

Verified Reactivity	Human, Mouse, Rat
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	This monoclonal antibody was raised against a fusion protein corresponding to amino acids 77-299 (PDZ domains 1 and 2) of human PSD-95.
Formulation	Phosphate-buffered solution.
Preparation	The antibody was purified by affinity chromatography.
Concentration	1 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Please note the storage condition for this antibody has been changed from -20°C to between 2°C and 8°C. You can also check your vial or your CoA to find the most accurate storage condition for this antibody.
Application	WB - Quality tested IHC-P - Verified
Recommended Usage	Each lot of this antibody is quality control tested by Western blotting . For Western blotting, the suggested use of this reagent is 0.1 - 5.0 µg/ml. For immunohistochemical staining of formalin-fixed paraffin-embedded tissue sections, a concentration range of 2.0 - 10 µg/ml is suggested. It is recommended that the reagent be titrated for optimal performance for each application.
Product Citations	1. Brown EA, <i>et al.</i> 2018. Mol Autism. 9:48. PubMed
RRID	AB_2734563 (BioLegend Cat. No. 810302) AB_2564749 (BioLegend Cat. No. 810301)

Antigen Details

Structure	Expected MW: 95-110 kD
Biology Area	Cell Biology, Neuroscience, Synaptic Biology
Molecular Family	Postsynaptic proteins
Gene ID	1742


Related Protocols

[Western Blotting Protocol](#)


Other Formats

Purified anti-PSD95


Product Data




Western blot of purified anti-PSD95 antibody (clone K28/74). Lane 1: Molecular weight marker; Lane 2: 10 μ g of human brain membrane lysate; Lane 3: 10 μ g of mouse brain membrane lysate; Lane 4: 10 μ g of rat brain membrane lysate. The blot was incubated with 0.5 μ g/mL of the primary antibody overnight at 4°C, followed by incubation with HRP labeled goat anti-mouse IgG (Cat. No. 405306). Enhanced chemiluminescence was used as the detection system.




IHC staining of purified anti-PSD95 antibody (clone K28/74) on formalin-fixed paraffin-embedded mouse brain tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 10 μ g/ml of the primary antibody overnight at 4°C. BioLegend's Ultra-Streptavidin (USA) HRP kit (Multi-Species, DAB, Cat. No. 929901) was used for detection followed by hematoxylin counterstaining, according to the protocol provided. The image was captured with a 40X objective. Scale bar: 50 μ m




IHC staining of purified anti-PSD95 antibody (clone K28/74) on formalin-fixed paraffin-embedded rat brain tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 10 μ g/ml of the primary antibody overnight at 4°C. BioLegend's Ultra-Streptavidin (USA) HRP kit (Multi-Species, DAB, Cat. No. 929901) was used for detection followed by hematoxylin counterstaining, according to the protocol provided. The image was captured with a 40X objective. Scale bar: 50 μ m



IHC staining of purified anti-PSD95 antibody (clone K28/74) on formalin-fixed paraffin-embedded human brain tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 10 $\mu\text{g/ml}$ of the primary antibody overnight at 4°C. BioLegend's Ultra-Streptavidin (USA) HRP kit (Multi-Species, DAB, Cat. No. 929901) was used for detection followed by hematoxylin counterstaining, according to the protocol provided. The image was captured with a 40X objective. Scale bar: 50 μm



IHC staining of purified anti-PSD95 antibody (clone K28/74) on formalin-fixed paraffin-embedded mouse cerebellum tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 10 $\mu\text{g/mL}$ of the primary antibody overnight at 4°C, followed by incubation with Alexa Fluor® 555 anti-mouse IgG for one hour at room temperature. Nuclei were counterstained with DAPI. The image was captured with a 40X objective. Scale bar: 50 μm



IHC staining of purified anti-PSD95 antibody (clone K28/74) on formalin-fixed paraffin-embedded mouse hippocampus tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 10 $\mu\text{g/mL}$ of the primary antibody overnight at 4°C, followed by incubation with Alexa Fluor® 555 anti-mouse IgG for one hour at room temperature. Nuclei were counterstained with DAPI. The image was captured with a 40X objective. Scale bar: 50 μm

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587