

Recombinant Mouse GM-CSF (carrier-free)

Catalog# / Size	576302 / 10 µg 576304 / 25 µg 576306 / 100 µg 576308 / 500 µg
Regulatory Status	RUO
Other Names	Granulocyte/macrophage-colony stimulating factor, CSF-α, Pluripoietin-α, Eosinophil colony stimulating factor (Eo-CSF), Burst promoting activity (BPA)
Description	GM-CSF is a hematopoietic factor that is produced by T cells, macrophages, fibroblasts and endothelial cells. This multifunctional cytokine stimulates progenitor cells of neutrophils, eosinophils and macrophages. GM-CSF is also a differentiation and activating factor for granulocytic and monocytic cells.

Product Details


Source	Mouse GM-CSF, amino acids Ala18-Lys141 (Accession # NM_009969) was expressed in <i>E. coli</i> .
Molecular Mass	The 125 amino acid N-terminal methionylated recombinant protein has a predicted molecular mass of 14.2 kDa. The DTT-reduced protein migrates at approximately 13 kDa and the non-reduced protein migrates at approximately 11kDa by SDS-PAGE.
Purity	Purity is >98%, as determined by Coomassie stained SDS-PAGE.
Formulation	0.22 µm filtered protein solution is in 10mM NaH ₂ PO ₄ , 150mM NaCl, pH 7.2.
Endotoxin Level	Endotoxin level is <0.1 EU/µg (<0.01ng/µg) protein as determined by the LAL method.
Concentration	10 and 25 µg sizes are bottled at 200 µg/mL. 100 µg size and larger sizes are lot-specific and bottled at the concentration indicated on the vial. To obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.
Storage & Handling	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to six months, or at -70°C or colder until the expiration date. For maximum results, quick spin vial prior to opening. The protein can be aliquoted and stored at -20°C or colder. Stock solutions can also be prepared at 50 - 100 µg/mL in appropriate sterile buffer, carrier protein such as 0.2 - 1% BSA or HSA can be added when preparing the stock solution. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated freeze/thaw cycles.
Activity	The ED ₅₀ is <0.05 ng/ml, corresponding to a specific activity >2x10 ⁷ units/mg.
Application	Bioassay
Recommended Usage	Use when high specific biological activity is required.
Application Notes	BioLegend carrier-free recombinant proteins provided in liquid format are shipped on blue-ice. Our comparison testing data indicates that when handled and stored as recommended, the liquid format has equal or better stability and shelf-life compared to commercially available lyophilized proteins after reconstitution. Our liquid proteins are verified in-house to maintain activity after shipping on blue ice and are backed by our 100% satisfaction guarantee . If you have any concerns, contact us at tech@biolegend.com .
Application References	1. Ahn J, <i>et al.</i> 2012. <i>PNAS</i> . 109:19386. PubMed
(PubMed link indicates BioLegend citation)	2. Verhagen J, <i>et al.</i> 2012. <i>PNAS</i> . 110:E221. PubMed
	3. Bretscher P, <i>et al.</i> 2015. <i>EMBO Mol Med</i> . 7:593. PubMed
Product Citations	1. Ayala M, <i>et al.</i> 2017. <i>J Cancer Res Clin Oncol</i> . 10.1007/s00432-017-2421-7. PubMed
	2. Ayala M, <i>et al.</i> 2017. <i>Breast Cancer Res Treat</i> . 10.1007/s10549-017-4414-2. PubMed
	3. Bachran C, <i>et al.</i> 2017. <i>Sci Rep</i> . 10.1038/s41598-017-17948-0. PubMed
	4. Nakamura K, <i>et al.</i> 2018. <i>Cancer Cell</i> . 33:634. PubMed
	5. Snell LM, <i>et al.</i> 2018. <i>Immunity</i> . 49:678. PubMed

6. Chen X *et al.* 2017. *Cell stem cell.* 21(6):747-760 . [PubMed](#)
7. Wang Y, *et al.* 2019. *Front Cell Infect Microbiol.* 9:286. [PubMed](#)
8. Paris J *et al.* 2019. *Cell Stem Cell.* 25(1):137-148 . [PubMed](#)
9. Xueyang Yu *et al.* 2017. *Immunity.* 47(5):903-912 . [PubMed](#)
10. Chowdhury D, *et al.* 2019. *Cell Rep.* 27:3873. [PubMed](#)
11. Li Y, *et al.* 2020. *Cell Rep.* 30:1753. [PubMed](#)
12. Gogokhia L, *et al.* 2019. *Cell Host Microbe.* 25:285. [PubMed](#)
13. Jain A, *et al.* 2020. *Nat Immunol.* 0.920138889. [PubMed](#)
14. Lajqi T, *et al.* 2020. *Front Immunol.* 11:546415. [PubMed](#)
15. Ahn J, *et al.* 2012. *Proc Natl Acad Sci U S A.* 109:19386. [PubMed](#)
16. Bretscher P, *et al.* 2015. *EMBO Mol Med.* 7:593. [PubMed](#)
17. Jin Z, *et al.* 2020. *NPJ Regen Med.* 5:19. [PubMed](#)
18. Markworth JF, *et al.* 2020. *JCI Insight.* 5:00. [PubMed](#)
19. Alban TJ, *et al.* 2020. *Front Immunol.* 1.285416667. [PubMed](#)
20. Delbridge ARD, *et al.* 2021. *Front Cell Neurosci.* 14:592005. [PubMed](#)


Antigen Details

Structure	cytokine
Distribution	T cells, monocytes/macrophages, fibroblasts, endothelial cells, mast cells
Function	Synergistic with IL-1, IL-3, G-CSF; E21R competitive antagonist for receptor binding; stored in ECM with heparan sulfate proteoglycans
Interaction	Granulocyte/macrophage/erythroid/megakaryocytic progenitors, myeloblasts, monoblasts
Ligand/Receptor	Heterodimer GM-CSFR μ subunit (CDw116); β -subunit (CDw131) in common with IL-3R, IL-5R
Bioactivity	Growth/development granulocyte/macrophage progenitors; differentiates myeloblasts/monoblasts; synergizes with Epo proliferation of erythroid/megakaryocytic progenitors
Cell Type	Embryonic Stem Cells, Hematopoietic stem and progenitors
Biology Area	Cell Biology, Stem Cells
Molecular Family	Cytokines/Chemokines, Growth Factors
Antigen References	<ol style="list-style-type: none"> 1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press San Diego. 2. Demetri G, <i>et al.</i> 1991. <i>Blood</i> 78:2791. 3. Fan D, <i>et al.</i> 1991. <i>In vivo</i> 5:571. 4. Negrin R, <i>et al.</i> 1992. <i>Adv. Pharmacol.</i> 23:263.
Gene ID	12981

Product Data



FDC-P1 cell proliferation induced by mouse GM-CSF.



Recombinant mouse GM-CSF induces the proliferation of mouse FDC-P1 cell line in a dose dependent manner. BioLegend's protein was compared side-by-side to the leading competitor's equivalent product.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587