

Biotin anti-Apo E, 154-162 Antibody

Catalog# / Size	852805 / 25 µg 852806 / 100 µg
Clone	A17065A
Regulatory Status	RUO
Other Names	Apolipoprotein E, APO-E, Alzheimer Disease 2, LDLQC5, APOE, LPG, AD2
Isotype	Mouse IgG2b, κ
Description	Apolipoprotein E (Apo E) belongs to a class of apolipoproteins that mediate the binding, internalization, and catabolism of lipoprotein particles. Apo E is produced by the liver and macrophages in peripheral tissues where it regulates cholesterol metabolism. In the central nervous system, Apo E is primarily produced by astrocytes and plays an important role in transporting cholesterol to neurons via Apo E receptors. Apo E has three alleles (Apo E ₂ , Apo E ₃ and Apo E ₄) that differ from each other by only one or two amino acids at positions 112 and 158. Apo E ₄ is one of the largest known genetic risk factors for late-onset Alzheimer's disease. Furthermore, Apo E is associated with cardiovascular disease, Lipoprotein Glomerulopathy and Sea-Blue Histiocyte Disease.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	KLH conjugated Apo E peptide encompassing amino acids 154-162.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	WB - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by Western blotting . For Western blotting, the suggested use of this reagent is 1.0 - 10 µg per ml. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	This antibody cross-reacts with Apo E ₂ , Apo E ₃ , and Apo E ₄ .
RRID	AB_2783425 (BioLegend Cat. No. 852805) AB_2783426 (BioLegend Cat. No. 852806)

Antigen Details

Structure	Human Apo E is a 299 amino acid protein with a molecular mass of 32 kD, and exists as 3 major alleles, Apo E ₂ , Apo E ₃ , and Apo E ₄ .
Distribution	Tissue sources: Ubiquitously expressed. Highest levels in liver, brain, spleen, adrenal gland, and kidney. Cellular distribution: Plasma membrane, nucleus, endoplasmic reticulum, golgi apparatus, endosome, and extracellular.
Function	Apo E is involved in lipoprotein metabolism as well as the transport of fat-soluble vitamins, and cholesterol. Apo E also forms complexes with amyloid beta peptides leading to its uptake and

degradation by microglia.

Interaction	Tau, amyloid beta
Ligand/Receptor	LDL receptor, SORL1.
Biology Area	Cell Biology, Neurodegeneration, Neuroscience, Protein Misfolding and Aggregation
Molecular Family	Apolipoproteins

Antigen References	<ol style="list-style-type: none">1. Mahley RW. 2016. <i>Arterioscler Thromb Vasc Biol.</i> 36: 1305-1315.2. Riedel BC, et al. 2016 <i>J Steroid Biochem Mol Biol.</i> 160:134-47.3. Tai LM, et al. 2016 <i>Acta Neuropathol.</i> 131(5):709-23.4. Kim J, et al. 2014. <i>Mol. Cells.</i> 37:7675. Holtzman DM, et al. 2012. <i>Cold Spring Harb. Perspect. Med.</i> 2:a006312
---------------------------	--

Gene ID [348](#)


Related Protocols

[Western Blotting Protocol](#)

Other Formats

Purified anti-Apo E, 154-162, Biotin anti-Apo E, 154-162, HRP anti-Apo E, 154-162

Product Data



Western blot of Biotin anti-Apo E, 154-162 antibody (clone A17065A). Lane 1: Molecular weight marker; Lane 2: 40 ng of Apo E2 recombinant human protein; Lane 3: 40 ng of Apo E3 recombinant human protein; Lane 4: 40 ng of Apo E4 recombinant human protein; Lane 5: 40 µg of human plasma. The blot was incubated with 1.0 µg/mL of the primary antibody overnight at 4°C, followed by incubation with HRP Streptavidin (Cat. No. 405210) for 60 minutes at room temperature. Enhanced chemiluminescence (Cat. No. 426302) was used as the detection system.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587