

PE/Dazzle™ 594 anti-human IL-4 Antibody

Catalog# / Size	500831 / 25 tests 500832 / 100 tests
Clone	MP4-25D2
Regulatory Status	RUO
Other Names	Interleukin-4, Ia inducing factor (IaIF), B-cell stimulating factor-1 (BSF-1), Hodgkin's cell growth factor (HCGF), Mast cell growth factor-2 (MCGF-2), Macrophage fusion factor (MFF), T cell growth factor-2 (TCGF-2)
Isotype	Rat IgG1, κ
Description	IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells, and basophils. IL-4 elicits many different biological responses but has two dominant functions. The first is regulating differentiation of naïve CD4 ⁺ T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated immune response controlled by Th1 cells. The second is regulating IgE and IgG1 production by B cells.

Product Details

Verified Reactivity	Human, Cross-Reactivity: Swine (Pig, Porcine), Rhesus
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	CHO-expressed, recombinant human IL-4
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. * PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	ELISA Detection^{1,3} or ELISPOT Detection^{4,5}: The biotinylated MP4-25D2 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified 8D4-8 antibody (Cat. No. 500702/500707) as the capture antibody. Flow Cytometry^{6,9}: The fluorochrome-labeled MP4-25D2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-4 -producing cells within mixed cell populations. Neutralization¹⁻³: The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of human IL-4 bioactivity (Cat. No. 500815). The MP4-25D2 antibody can neutralize the bioactivity of natural or recombinant IL-4.
Application References	1. Chretien I, <i>et al.</i> 1989. <i>J. Immunol. Methods</i> 117:67. (ELISA Detection, Neut) 2. Ramanathan L, <i>et al.</i> 1993. <i>Biochem.</i> 32:3549. (Neut) 3. Abrams J, <i>et al.</i> 1992. <i>Immunol. Rev.</i> 127:5. (ELISA Detection, Neut)
(PubMed link indicates BioLegend citation)	

4. Mahanty S, *et al.* 1992. *J. Immunol.* 148:3567. (ELISPOT Detection)
5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19. (ELISPOT Detection)
6. Prussin C, *et al.* 1995. *J. Immunol. Methods* 188:117. (ICFC)
7. Raqib R, *et al.* 1995. *Infect. Immun.* 63:289.
8. Andersson J, *et al.* 1994. *Immunology* 83:16.
9. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
10. Kubota M, *et al.* 1997. *J. Immunol.* 158:5321.
11. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
12. Kroneke MA, *et al.* 2012. *J. Immunol.* 188:3734. [PubMed](#)

Product Citations

1. Thieme CJ, *et al.* 2020. *Cell Rep Med.* 1:100092. [PubMed](#)

RRID

AB_2564035 (BioLegend Cat. No. 500831)
 AB_2564036 (BioLegend Cat. No. 500832)

Antigen Details

Structure	Cytokine; 15-19 kD (Mammalian)
Bioactivity	Differentiation of naïve CD4 ⁺ T cells to the T _H 2 type, proliferation/differentiation of activated B cells, expression of class II MHC antigens, and of low affinity IgE receptors in resting B cells
Cell Sources	Mast cells, T cells, bone marrow stromal cells
Cell Targets	B cells, T cells, monocytes, endothelial cells, fibroblasts
Receptors	Heterodimer IL-4R α (CD124); γ -subunit (CD132) in common with IL-2R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none"> 1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook</i>. Academic Press San Diego. 2. Boulay J, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:294. 3. Dullens H, <i>et al.</i> 1991. <i>In vivo</i> 5:567. 4. Paul W. 1991. <i>Blood</i> 77:1859.
Regulation	Upregulated by IL-2, platelet activating factor; downregulated by TGF- β
Gene ID	3565

Related Protocols



[Intracellular Cytokine Staining Protocol - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

PE/Dazzle™ 594 anti-human IL-4, APC/Cyanine7 anti-human IL-4, Brilliant Violet 510™ anti-human IL-4, Ultra-LEAF™ Purified anti-human IL-4, Alexa Fluor® 488 anti-human IL-4, Alexa Fluor® 647 anti-human IL-4, PerCP/Cyanine5.5 anti-human IL-4, PE/Cyanine7 anti-human IL-4, Brilliant Violet 421™ anti-human IL-4, Brilliant Violet 605™ anti-human IL-4, Purified anti-human IL-4 (Maxpar® Ready), APC anti-human IL-4, Biotin anti-human IL-4, FITC anti-human IL-4, PE anti-human IL-4, Purified anti-human IL-4

Product Data



PMA + Ionomycin-stimulated (6 hours) human peripheral blood lymphocytes (in the presence of monensin) were stained with CD3 FITC, fixed, permeabilized, and then stained with IL-4 (clone MP4-25D2) PE/Dazzle™ 594 (top) or rat IgG1, κ PE/Dazzle™ 594 isotype control (bottom).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
 Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587