

APC anti-human HLA-DR Antibody

Catalog# / Size	327021 / 25 tests 327022 / 100 tests
Clone	LN3
Regulatory Status	RUO
Other Names	Major Histocompatibility Class II, MHC class II
Isotype	Mouse IgG2b, κ
Description	The LN3 monoclonal antibody reacts with the HLA-DR antigen, a member of MHC class II molecules. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4 ⁺ T cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	human PBL
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining ¹ of frozen sections and formalin-fixed paraffin-embedded sections ¹ , and immunoprecipitation ¹ .
Application References	<ol style="list-style-type: none"> 1. Marder RJ, <i>et al.</i> 1985. <i>Lab. Invest.</i> 52:497. 2. Norton AJ and Isaacson PG. 1987. <i>Am. J. Pathol.</i> 128:225. 3. Hua ZX, <i>et al.</i> 1998. <i>Hum. Pathol.</i> 29(12):1441.
RRID	AB_2734310 (BioLegend Cat. No. 327021) AB_2734311 (BioLegend Cat. No. 327022)

Antigen Details

Structure	Ig superfamily, MHC class II, heterodimeric transmembrane protein
------------------	---

Distribution	B cells, activated T cells, monocytes/macrophages, dendritic cells, other APCs
Function	Peptide presentation
Ligand/Receptor	CD3/TCR, CD4
Cell Type	Antigen-presenting cells, B cells, Dendritic cells, Macrophages, Monocytes, T cells, Tregs
Biology Area	Immunology, Innate Immunity
Molecular Family	MHC Antigens
Antigen References	<ol style="list-style-type: none"> 1. Levacher M, <i>et al.</i> 1990. <i>Clin. Exp. Immunol.</i> 81:177. 2. Terstappen L, <i>et al.</i> 1990. <i>J. Leuk. Biol.</i> 48:138. 3. Edwards J, <i>et al.</i> 1985. <i>J. Immunol.</i> 137:490. 4. van Es A, <i>et al.</i> 1984. <i>Transplantation</i> 37:65. 5. O'Doherty U, <i>et al.</i> 1994. <i>Immunology</i> 82:487. 6. Thomas R, <i>et al.</i> 1994. <i>J. Immunol.</i> 153:4016. 7. Grouard G, <i>et al.</i> 1996. <i>Nature</i> 384:364.
Gene ID	3122 3123


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human HLA-DR, Biotin anti-human HLA-DR, FITC anti-human HLA-DR, PE anti-human HLA-DR, Alexa Fluor® 488 anti-human HLA-DR, Alexa Fluor® 647 anti-human HLA-DR, Alexa Fluor® 700 anti-human HLA-DR, Pacific Blue™ anti-human HLA-DR, PE/Cyanine7 anti-human HLA-DR, PerCP/Cyanine5.5 anti-human HLA-DR, APC anti-human HLA-DR, APC/Fire™ 750 anti-human HLA-DR

Product Data



Human peripheral blood lymphocytes were stained with HLA-DR (clone LN3) APC (filled histogram) or mouse IgG2b, κ APC isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

