

PerCP/Cyanine5.5 anti-Histone H3-Phosphorylated (Ser28) Antibody

Catalog# / Size	641013 / 25 tests 641014 / 100 tests
Clone	HTA28
Regulatory Status	RUO
Other Names	Histone-H3
Isotype	Rat IgG2a, κ
Description	H3 is a core component of the nucleosome that serves to wrap and compact DNA into chromatin. Histones therefore, limit the accessibility of DNA, providing mechanisms for transcription regulation, DNA repair and replication and chromosomal stability. During mitosis, H3 is phosphorylated at serine 28. This phosphorylation coincides with chromosome condensation initiated at prophase and disappears at late anaphase. H3 has been demonstrated to be phosphorylated by the action of MLTK- α (mixed lineage kinase-like mitogen activated protein triple kinase α) in response to ultraviolet B light and epidermal growth factor, as well as Aurora-B during mitosis.

Product Details

Reactivity	Human
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Synthetic peptide conjugated to KLH, corresponding to amino acids 23-35 of human histone H3.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood. * PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
Application Notes	This clone is not recommended for ChIP (Chromatin Immunoprecipitation) assays (as determined by in-house testing).
Additional Product Notes	BioLegend is in the process of converting the name PerCP/Cy5.5 to PerCP/Cyanine5.5. The dye molecule remains the same, so you should expect the same quality and performance from our PerCP/Cyanine5.5 products. Contact Technical Service if you have any questions.
Application References	<ol style="list-style-type: none"> Hirata A, et al. 2004. <i>J. Histochem. Cytochem.</i> 52:1503. Goto H, et al. 1999. <i>J. Biol. Chem.</i> 274:25543. Ozawa K. 2008. <i>Cytometry A</i> 73:517. Goode NJ, et al. 2014. <i>PLoS Genet.</i> 10:1004323. PubMed
RRID	AB_2565891 (BioLegend Cat. No. 641013) AB_2565892 (BioLegend Cat. No. 641014)

Antigen Details

Structure	H3 is part of the nucleosome, comprised of an octameric complex with H2A, H2B, and H4 proteins.
Distribution	Nucleus
Function	H3 is a core component of the nucleosome that serves to wrap and compact DNA into chromatin. Histones therefore, limit the accessibility of DNA, providing mechanisms for transcription regulation, DNA repair and replication and chromosomal stability.
Interaction	Two molecules of H3 form a heterotetramer with two molecules of H4.
Biology Area	Cell Biology, DNA Repair/Replication, Transcription Factors
Molecular Family	Phospho-Proteins
Antigen References	1. Choi HS, <i>et al.</i> 2005. <i>J. Biol. Chem.</i> 280:13545. 2. Goto H, <i>et al.</i> 2002. <i>Genes Cells</i> 7:11. 3. Garcia BA, <i>et al.</i> 2005. <i>Biochemistry</i> 44:13202.
Regulation	H3 is regulated by acetylation, methylation, citrullination, phosphorylation, and ubiquitination.
Gene ID	8290

Related Protocols


[Intracellular Cytokine Staining Protocol - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-Histone H3-Phosphorylated (Ser28), Alexa Fluor® 488 anti-Histone H3-Phosphorylated (Ser28), Alexa Fluor® 647 anti-Histone H3-Phosphorylated (Ser28), Purified anti-Histone H3-Phosphorylated (Ser28) (Maxpar® Ready), PE anti-Histone H3-Phosphorylated (Ser28), PE/Cyanine7 anti-Histone H3-Phosphorylated (Ser28), PerCP/Cyanine5.5 anti-Histone H3-Phosphorylated (Ser28), Alexa Fluor® 594 anti-Histone H3-Phosphorylated (Ser28), Direct-Blot™ HRP anti-Histone H3 Phospho (Ser28)

Product Data



HeLa cells untreated (open histogram), and HeLa cells treated with Nocodazole for 24-hours (filled histogram), were fixed and permeabilized with 70% ethanol and then stained with anti-Histone H3-Phosphorylated (Ser28) (clone HTA28) PerCP/Cyanine5.5.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587

