

## APC anti-human CD86 Antibody

<b>Catalog# / Size</b>	374207 / 25 tests 374208 / 100 tests
<b>Clone</b>	BU63
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	HCDM listed
<b>Other Names</b>	B7-2, B70, Ly-58
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, monocytes/macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is the ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce costimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can bind to CD152 as well, also known as CTLA-4, to deliver an inhibitory signal to T cells.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	ARH 77 (B lymphoblastoid cell line).
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ l per million cells in 100 $\mu$ l staining volume or 5 $\mu$ l per 100 $\mu$ l of whole blood.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Product Citations</b>	1. Dinh HQ, <i>et al.</i> 2020. Immunity. 53(2):319-334.e6. <a href="#">PubMed</a>
<b>RRID</b>	AB_2721448 (BioLegend Cat. No. 374207) AB_2721449 (BioLegend Cat. No. 374208)

### Antigen Details

---

<b>Structure</b>	Ig superfamily, single-chain transmembrane glycoprotein, 80 kD
<b>Distribution</b>	Monocytes/macrophages, activated B cells and T cells, dendritic cells

<b>Function</b>	T Cell activation
<b>Interaction</b>	C-Jun, NFκB
<b>Ligand/Receptor</b>	CD28, CD152
<b>Cell Type</b>	B cells, Dendritic cells, Macrophages, Monocytes, T cells
<b>Biology Area</b>	Cell Biology, Costimulatory Molecules, Immunology, Neuroscience, Neuroscience Cell Markers
<b>Molecular Family</b>	CD Molecules, Immune Checkpoint Receptors
<b>Antigen References</b>	1. Hathcock K, <i>et al.</i> 1996. <i>Adv. Immunol.</i> 62:131. 2. June C, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:321.
<b>Gene ID</b>	<a href="#">942</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

Purified anti-human CD86, FITC anti-human CD86, PE anti-human CD86, APC anti-human CD86, PE/Cyanine7 anti-human CD86, Brilliant Violet 421™ anti-human CD86, Brilliant Violet 605™ anti-human CD86, PerCP/Cyanine5.5 anti-human CD86, PE/Dazzle™ 594 anti-human CD86

## Product Data



Human peripheral blood monocytes stained with CD14 PerCP/Cy5.5 and human CD86 (clone BU63, top) APC or Mouse IgG1, κ APC isotype control (bottom).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic,

therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587