

Purified anti-Tau, 185-195 Antibody (Previously Covance catalog# MMS-5018)

Catalog# / Size	814402 / 25 µL 814401 / 100 µL
Clone	77E9
Regulatory Status	RUO
Other Names	Microtubule-associated protein tau, PHF-tau, Paired Helical Filament tau
Previously	Covance Catalog# MMS-5018
Isotype	Mouse IgG1, κ
Description	<p>Tau proteins are microtubule-associated protein (MAPs) which are abundant in neurons of the central nervous system, but are also expressed at very low levels in CNS astrocytes and oligodendrocytes and elsewhere. One of tau's main functions is to modulate the stability of axonal microtubules. Tau is active primarily in the distal portions of axons providing microtubule stabilization as well as flexibility. Pathologies and dementias of the nervous system such as Alzheimer's disease feature tau proteins that have become defective and no longer stabilize microtubules properly. As a result, tau forms aggregates with specific structural properties referred to as Paired Helical Filaments (PHFs) that are a characteristic of many different types of dementias, known as tauopathies.</p> <p>Tau has two primary ways of controlling microtubule stability: isoforms and phosphorylation. Six tau isoforms exist in human brain tissue, and they are distinguished by the number of binding domains. Three isoforms have three binding domains and the remaining three have four binding domains. The binding domains are located in the carboxy-terminus of the protein and are positively-charged (for binding to the negatively-charged microtubule). Tau isoforms with four binding domains are better at stabilizing microtubules than those with three binding domains.</p> <p>Thus, in the human brain, the tau proteins constitute a family of six isoforms with the range from 352-441 amino acids. They also differ in either zero, one or two inserts of 29 amino acids at the N-terminal part (exon 2 and 3), and three or four repeat-binding regions at the C-terminus. So, the longest isoform in the CNS has four repeats (R1, R2, R3 and R4) and two inserts (441 amino acids total), while the shortest isoform has three repeats (R1, R3 and R4) and no insert (352 amino acids total). Tau is also a phosphoprotein with 79 potential Serine (Ser) and Threonine (Thr) phosphorylation sites on the longest tau isoform. Phosphorylation has been reported on approximately 30 of these sites in normal tau proteins. Mechanisms that drive tau lesion formation in the highly prevalent sporadic form of AD are not fully understood, but appear to involve abnormal post-translational modifications (PTMs) that influence tau function, stability, and aggregation propensity.</p>

Product Details

Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	This monoclonal antibody was raised against full-length human tau.
Formulation	Phosphate-buffered solution.
Preparation	The antibody was purified by affinity chromatography.
Concentration	1 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Please note the storage condition for this antibody has been changed from -20°C to between 2°C and 8°C. You can also check your vial or your CoA to find the most accurate storage condition for this antibody.
Application	WB - Quality tested IHC-P - Verified

Recommended Usage

Each lot of this antibody is quality control tested by [Western blotting](#). For Western blotting, the suggested use of this reagent is 1.0 - 10 µg per ml. For immunohistochemistry on formalin-fixed paraffin-embedded tissue sections, a concentration range of 0.5 - 1.0 µg/ml is suggested. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes

BioLegend is the Exclusive Provider of this Antibody Product

Product Citations

1. Vos A, *et al.* 2015. *Alzheimers Dement.* 11: 1461-1469. [PubMed](#)

RRID

AB_2715845 (BioLegend Cat. No. 814402)
AB_2564783 (BioLegend Cat. No. 814401)

Antigen Details

Biology Area

Cell Biology, Neurodegeneration, Neuroscience, Protein Misfolding and Aggregation

Molecular Family

Tau

Gene ID

[4137](#)


Related Protocols

[Western Blotting Protocol](#)


Other Formats

Purified anti-Tau, 185-195

Product Data



Western blot of purified anti-Tau, 185-195 antibody (clone 77E9). Lane 1: Molecular weight marker; Lane 2: 5 µl of recombinant human Tau protein ladder; Lane 3: 20 µg of normal human brain lysate; Lane 4: 20 µg of human Alzheimer's disease brain lysate. The blot was incubated with 1 µg/mL of the primary antibody overnight at 4°C, followed by incubation with HRP labeled goat anti-mouse IgG (Cat. No. 405306). Enhanced chemiluminescence was used as the detection system.



IHC staining of purified anti-Tau, 185-195 antibody (clone 77E9) on formalin-fixed paraffin-embedded human Alzheimer's disease brain tissue. Following antigen retrieval using Sodium Citrate H.I.E.R., the tissue was incubated with 1 μ g/ml of the primary antibody for 20 minutes at room temperature. BioLegend's Ultra-Streptavidin (USA) HRP kit (Multi-Species, DAB, Cat. No. 929901) was used for detection followed by hematoxylin counterstaining, according to the protocol provided. The image was captured with a 40X objective. Scale bar: 50 μ m

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587